

Benha University Faculty of Engineering- Shoubra Eng. Mathematics & Physics Department Preparatory Year		Final Term Exam Date: December 16, 2017 Course: Mathematics 1 – A Duration: 3 hours
- The Exam consists of one page - Answer All Questions	(تخلفات)	- No. of questions: 4 - Total Mark: 100
Question 1		
Find y' from the following:		
(a) $y = 2x^4 + 4^x + x$	(b) $y = (x - \cos x)^6$	(c) $y = \log x - \ln(1 + \sin x)$
(d) $y = \tan^2 x + \sin x^3$	(e) $y^4 = x \ln y + \sin x$	(f) $y = t \cos t, x = t - \ln t$
Question 2		
(a) Find the following limits:		
(i) $\lim_{x \rightarrow 1} \frac{x^7 - 1}{1 - x^5}$	(ii) $\lim_{x \rightarrow 0} \frac{\sin 2x}{2^x - 3^x}$	(iii) $\lim_{x \rightarrow 0} \frac{x - \sin x}{x^3 + 2x}$
(iv) $\lim_{x \rightarrow \infty} \frac{x^4 + 2x}{x + 3x^5}$		
(b) Write the Maclurin's expansion of the function : $f(x) = (x + 1) \sin x$.		
(c) Sketch the curve of the function : $f(x) = x + \frac{1}{x}$		
(d) State and verify Rolle's theorem for, $f(x) = (x - 1)^2$ in the interval $[0, 2]$.		
(e) Find the integrals: (i) $\int (x^3 + 3^x) dx$ (ii) $\int (1 - 2^x)^2 dx$ (iii) $\int (1 + \cos 2x) dx$		
Question 3		
(a) Prove that : $1 + 2 + 3 + \dots n = \frac{1}{2}n(n + 1)$ for $n \geq 1$.		
(b) Find the sum : $\sum_{r=1}^n (4r^2 - 1)$		
(c) Find the coefficient of x^{15} in the expansion : $(1 + x)^4 \cdot (1 - x)^{-4}$		
(d) Express $\cos 3\theta$ and $\sin 3\theta$ in terms of $\cos \theta$ and $\sin \theta$.		
Question 4		
(a) Resolve the fraction : $\frac{x^4 - 2x^3 + 3x^2 - x + 3}{x^3 - 2x^2 + 3x}$ into its partial fractions.		
(b) Write the first four terms in the expansion : $\sqrt[3]{8 - 2x}$.		
(c) Use the Gauss-Jordan algorithm to solve the linear system :		
$x_1 - 2x_2 + 2x_3 = 2, 3x_1 + x_2 - x_3 = 6, 2x_1 - 2x_2 - 3x_3 = -6$		
(d) Show that $x = 2i$ is a zero to the polynomial		
$f(x) = x^4 - 5x^3 + 10x^2 - 20x + 24$ and then solve the equation $f(x) = 0$.		

Good Luck

Dr. Mohamed Eid

Dr. Fathi Abdallah

Good Luck,

Dr. Mohamed Eid

Dr. Fathi Abdsallam

Model Answer

Answer of Question 1

$$(a) y' = 8x^3 + 4^x \cdot \ln 4 + 1$$

$$(b) y' = 6(x - \cos x)^5 (1 + \sin x)$$

$$(c) y' = \frac{1}{\ln 10} \cdot \frac{1}{x} - \frac{\cos x}{1 + \sin x}$$

$$(d) y' = 2 \tan x \cdot \sec^2 x + \cos x^3 \cdot 3x^2$$

$$(e) 4y^3 \cdot y' = \ln y + x \cdot \frac{y'}{y} + \cos x$$

$$(f) y' = \frac{\cos t - t \cdot \sin t}{1 - \frac{1}{t}}$$

-----24-Marks

Answer of Question 2

$$(a)(i) \lim_{x \rightarrow 1} \frac{x^7 - 1}{1 - x^5} = \frac{0}{0} = -\frac{7}{5}$$

$$(ii) \lim_{x \rightarrow 0} \frac{\sin 2x}{2^x - 3^x} = \frac{0}{0} = \frac{2}{\ln(2/3)}$$

$$(iii) \lim_{x \rightarrow 0} \frac{x - \sin x}{x^3 + 2x} = \frac{0}{0} = \lim_{x \rightarrow 0} \frac{1 - \cos x}{3x^2 + 2} = \frac{0}{2} = 0$$

$$(iv) \lim_{x \rightarrow \infty} \frac{x^4 + 2x}{x + 3x^5} = \frac{\infty}{\infty} = \lim_{x \rightarrow \infty} \frac{1 + \frac{2}{x^3}}{\frac{1}{x^3} + 3x} = \frac{1}{\infty} = 0$$

-----8-Marks

$$(b) f(x) = (x + 1) \sin x = 0 + x + 2x^2 - \frac{1}{6}x^3 \dots$$

-----4-Marks

(c) The domain of this function is $\mathbb{R} - \{0\}$.

This function is odd.

The curve does not intersect y-axis.

At $y = 0$, we get the equation $x + \frac{1}{x} \neq 0$ which has no real roots. Then the curve does not intersect x-axis.

From $f'(x) = 1 - \frac{1}{x^2} = 0$, we get the equation $x^2 - 1 = 0$

which has the roots $x = 1, -1$.

Since $f''(x) = \frac{2}{x^3}$.

Since $f''(1) = 2$ is positive, then $(1, f(1)) = (1, 2)$ is minimum point.


Since $f''(-1) = -2$ is negative, then $(-1, f(-1)) = (-1, -2)$ is maximum point.

This function has no inflection points because $f''(x) = \frac{2}{x^3} \neq 0$.

This function has vertical asymptotic line $x = 0$ and has no horizontal asymptote because

$$\lim_{x \rightarrow \infty} f(x) = \infty.$$

It has inclined asymptotic line $y = x$.



The graph of f is shown in the Figure.

-----4-Marks

(d) We see that $f(x)$, its derivative $f'(x) = 2(x - 1)$ are continuous in the given interval $[0, 2]$ and $f(0) = f(2) = 1$.

Then $f'(c) = 2(c - 1) = 0$. Hence $c = 1$.

-----4-Marks

$$(e)(i) \int (x^3 + 3^x) dx = \frac{1}{4}x^4 - \frac{3^x}{\ln 3} + c$$

$$(ii) \int (1 - 2^x)^2 dx = \int (1 - 2 \cdot 2^x + 4^x) dx = x + 2 \frac{2^x}{\ln 2} + \frac{4^x}{\ln 4} + c$$

-----6-Marks

Dr. Mohamed Eid

Answer of Question 3

(a) Prove that $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$ for any integer $n \geq 1$.

Proof:

STEP 1: For $n=1$ $P(1)$ is true, since $1 = 1(1+1)/2$.

STEP 2: Suppose $P(k)$ is true for some $k \geq 1$,

$$\text{That is} \quad 1 + 2 + 3 + \dots + k = \frac{k(k+1)}{2}.$$

STEP 3: Prove that $P(k+1)$ is true for $n=k+1$,

$$\text{That is} \quad 1 + 2 + 3 + \dots + k + (k+1) = \frac{(k+1)(k+2)}{2}.$$

$$\text{We have} \quad 1 + 2 + 3 + \dots + k + (k+1) = \frac{k(k+1)}{2} + (k+1) = \frac{(k+1)(k+2)}{2},$$

Which show that $P(k+1)$ is true

-----6-Marks

(b) Find the sum $\sum_{r=1}^n (4r^2 - 1)$

$$\sum_{r=1}^n (4r^2 - 1) = \sum_{r=1}^n (2r-1)(2r+1)$$

$$u_r = (2r-1)(2r+1)$$

$$= \frac{(2r-1)(2r+1)(2r+3)}{6} - \frac{(2r-3)(2r-1)(2r+1)}{6}$$

$$= f(r+1) - f(r)$$

$$S_n = \frac{(2n-1)(2n+1)(2n+3)}{6} + \frac{1}{2}$$

$$\text{then} \quad \boxed{\sum_{r=1}^n (4r^2 - 1) = \frac{(2n-1)(2n+1)(2n+3)}{6} + \frac{1}{2}}$$

-----6-Marks

(c) Find coefficient x^{15} in the expansion $(1+x)^4(1-x)^{-4}$.

$$\begin{aligned} (1+x)^4(1-x)^{-4} &= \left(1 + 4x + 6x^2 + 4x^3 + x^4\right) \left(\sum_{r=0}^{\infty} {}^{-4}C_r (-x)^r\right) \\ &= \left(1 + 4x + 6x^2 + 4x^3 + x^4\right) \left(\sum_{r=0}^{\infty} (-1)^r {}^4C_r {}^{4+r-1}(-1)^r x^r\right) \end{aligned}$$

$$\begin{aligned}
&= (1 + 4x + 6x^2 + 4x^3 + x^4) \left(\sum_{r=0}^{\infty} C_r^{4+r-1} x^r \right) \\
&= (1 + 4x + 6x^2 + 4x^3 + x^4) \left(C_0^3 + C_1^4 x + C_2^5 x^2 + \dots + C_n^{3+n} x^n + \dots \right)
\end{aligned}$$

Coefficient x^{15} is

$$\begin{aligned}
&C_{15}^{18} + 4C_{14}^{17} + 6C_{13}^{16} + 4C_{12}^{15} + C_{11}^{14} \\
&= C_3^{18} + 4C_3^{17} + 6C_3^{16} + 4C_3^{15} + C_3^{14} \\
&= \frac{18.17.16}{3.2.1} + 4 \cdot \frac{17.16.15}{3.2.1} + 6 \cdot \frac{16.15.14}{3.2.1} + 4 \cdot \frac{15.14.13}{3.2.1} + \frac{14.13.12}{3.2.1} = 9080
\end{aligned}$$

-----6-Marks

(d) Express $\cos 3\theta$ and $\sin 3\theta$ in terms of $\cos \theta$ and $\sin \theta$

Using the equation

$$\begin{aligned}
(\cos n\theta + i \sin n\theta) &= (\cos \theta + i \sin \theta)^3 = \\
&= \cos^3 \theta + 3\cos^2 \theta (i \sin \theta) + 3\cos \theta (i \sin \theta)^2 + (i \sin \theta)^3 \\
&= \cos^3 \theta + 3i \cos^2 \theta (\sin \theta) - 3\cos \theta \sin^2 \theta - i \sin^3 \theta \\
&= \cos^3 \theta - 3\cos \theta \sin^2 \theta + i (3\cos^2 \theta \sin \theta - \sin^3 \theta)
\end{aligned}$$

Equating real and imaginary parts gives

$$\cos 3\theta = \cos^3 \theta - 3\cos \theta \sin^2 \theta$$

$$\sin 3\theta = 3\cos^2 \theta \sin \theta - \sin^3 \theta$$

-----6-Marks

Answer of Question 4

$$(a) \frac{x^4 - 2x^3 + 3x^2 - x + 3}{x^3 - 2x^2 + 3x}$$

$$\frac{x^4 - 2x^3 + 3x^2 - x + 3}{x^3 - 2x^2 + 3x} = x + \frac{-x + 3}{x(x^2 - 2x + 3)}$$

$$\frac{-x + 3}{x(x^2 - 2x + 3)} = \frac{A}{x} + \frac{Bx + C}{x^2 - 2x + 3}$$

-----6-Marks

$$(b) \sqrt[3]{8 - 2x}$$

$$\sqrt[3]{8-2x} = 2\left(1 - \frac{x}{4}\right)^{\frac{1}{3}} = 2\left[1 - \frac{1}{3}\left(\frac{x}{4}\right) + \frac{1}{2!}\left(\frac{1}{3}\right)\left(\frac{1}{3}-1\right)\left(\frac{x}{4}\right)^2 - \frac{1}{3!}\left(\frac{1}{3}\right)\left(\frac{1}{3}-1\right)\left(\frac{1}{3}-2\right)\left(\frac{x}{4}\right)^3 + \dots\right]$$

$$= 2\left[1 - \frac{x}{12} - \frac{x^2}{144} - \frac{5x^3}{5184} + \dots\right]$$

-----6-Marks

(c) Use the Gauss-Jordan algorithm to solve the linear system :

$$x_1 - 2x_2 + 2x_3 = 2, \quad 3x_1 + x_2 - x_3 = 6, \quad 2x_1 - 2x_2 - 3x_3 = -6$$

Solution

The augmented matrix of the system is

$$\left(\begin{array}{ccc|c} 1 & -2 & 2 & 2 \\ 3 & 1 & -1 & 6 \\ 2 & -2 & -3 & -6 \end{array}\right)$$

Applying Gauss-Jordan algorithm to this matrix yields

$$\left(\begin{array}{ccc|c} 1 & -2 & 2 & 2 \\ 0 & 7 & -7 & 0 \\ 0 & 2 & -7 & -10 \end{array}\right) \sim \left(\begin{array}{ccc|c} 1 & -2 & 2 & 2 \\ 0 & 1 & -1 & 0 \\ 0 & 2 & -7 & -10 \end{array}\right)$$

$$\sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & -5 & -10 \end{array}\right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 2 \end{array}\right) \sim \left(\begin{array}{ccc|c} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 2 \end{array}\right)$$

the system has unique solution $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix}$

-----7-Marks

(d) Show that $x = 2i$ is a zero to the polynomial

$$f(x) = x^4 - 5x^3 + 10x^2 - 20x + 24 \text{ and then solve the equation } f(x) = 0.$$

Solution

Divide the given polynomial by the linear expression $(x - 2i)$ and we found that the remainder $R = f(x - 2i) = 0$

this means that the value $2i$ is a zero of $f(x)$ and $2i$ is a root for $f(x) = 0$, then $-2i$ is another root and

$$f(x) = (x - 2i)(x + 2i)Q(x)$$

$$= (x^2 + 4)Q(x)$$

The polynomial $Q(x)$ is obtained by synthetic division as follows

2i		1	-5	10	-20	24	
			2i	-4-10i	20+12i	-24	
-2i		1	-5+2i	6-10i	12i		0
			-2i	10i	12i		
		1	-5	6		0	

Then

$$f(x) = (x^2 + 4)(x^2 - 5x + 6) = (x^2 + 4)(x - 2)(x - 3)$$

The roots are $\pm 2i$, 2, 3

-----7-Marks

Dr. Fathi Abdsallam